


# Introduction to Logic

## *Equality*


Michael Genesereth  
Computer Science Department  
Stanford University

# Isomorphic Representation

People





Arithmetic



# Homomorphic Representation

People (Nicknames):

*michael*  $\leftrightarrow$    
*mike*  $\leftrightarrow$  

Arithmetic:

*plus*(*s*(0), *s*(0))  $\leftrightarrow$   
*times*(*s*(*s*(0)), *s*(0))  $\leftrightarrow$  2  
*s*(*s*(0))  $\leftrightarrow$

# Two Approaches

## Approach 1 - Equality:

*mike = michael*

*father(michael) = william*

*Equivalence of terms in Herbrand universe*

---

## Approach 2 - Evaluable Terms (nicknames):

*s(0) + s(0) --> s(s(0))*

*s(s(0)) \* s(0) --> s(s(0))*

*Add new terms (terms not in the Herbrand universe)  
that "evaluate" to terms in Herbrand universe*

Equality

# Partitioning the Herbrand Universe

People:


<i>michael</i> <i>mike</i>	<i>maureen</i>	<i>katherine</i> <i>kathy</i> <i>kate</i>	...
-------------------------------	----------------	---	-----

Arithmetic:

...	<i>plus(s(0), s(0))</i> <i>times(s(s(0)), s(0))</i> <i>s(s(0))</i>	...
-----	--	-----

# Equality

Co-referential terms:



*plus(s(0), s(0))*  
*times(s(s(0)), s(0))*  
*s(s(0))*

Equality:

*equal(plus(s(0),s(0)), s(s(0)))*  
*equal(times(s(s(0)), s(0)), s(s(0)))*

# Syntactic Sugar

Formal Syntax:

*equal(plus(s(0),s(0)), s(s(0)))*

*equal(times(s(s(0)),s(0)), s(s(0)))*

Syntactic Sugar:

$$s(0) + s(0) = s(s(0))$$

$$s(s(0)) * s(0) = s(s(0))$$



# What is Needed

(1) Axioms that define properties of equality.

*michael = mike*

*mike = michael*

(2) Axioms that ensure that any true sentence that mentions a given term is also true of the sentence in which that term is replaced by an equivalent term.

*older(michael, maureen) ⇔ older(mike, maureen)*

# Equality Axioms

Reflexivity

$$\forall x.(x=x)$$

Symmetry

$$\forall x.\forall y.(x=y \Rightarrow y=x)$$

Transitivity

$$\forall x.\forall y.\forall z.(x=y \wedge y=z \Rightarrow x=z)$$

# Equality Proof

1.  $b = a$  Premise
2.  $b = c$  Premise
3.  $\forall x.(x = x)$  Premise
4.  $\forall x.\forall y.(x = y \Rightarrow y = x)$  Premise
5.  $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$  Premise

# Equality Proof

- |    |  |           |
|----|--|-----------|
| 1. | $b = a$  | Premise   |
| 2. | $b = c$  | Premise   |
| 3. | $\forall x.(x = x)$  | Premise   |
| 4. | $\forall x.\forall y.(x = y \Rightarrow y = x)$                        | Premise   |
| 5. | $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$ | Premise   |
| 6. | $b = a \Rightarrow a = b$  | 2 x UE: 4 |
| 7. | $a = b$  | IE: 6, 1  |

# Equality Proof

- |    |  |           |
|----|--|-----------|
| 1. | $b = a$  | Premise   |
| 2. | $b = c$  | Premise   |
| 3. | $\forall x.(x = x)$  | Premise   |
| 4. | $\forall x.\forall y.(x = y \Rightarrow y = x)$                        | Premise   |
| 5. | $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$ | Premise   |
| 6. | $b = a \Rightarrow a = b$  | 2 x UE: 4 |
| 7. | $a = b$  | IE: 6, 1  |
| 8. | $a = b \wedge b = c \Rightarrow a = c$                                 | 3 x UE: 5 |

# Equality Proof

1.	$b = a$	Premise
2.	$b = c$	Premise
3.	$\forall x.(x = x)$	Premise
4.	$\forall x.\forall y.(x = y \Rightarrow y = x)$	Premise
5.	$\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$	Premise
6.	$b = a \Rightarrow a = b$	2 x UE: 4
7.	$a = b$	IE: 6, 1
8.	$a = b \wedge b = c \Rightarrow a = c$	3 x UE: 5
9.	$a = b \wedge b = c$	AI: 7, 2
10.	$a = c$	IE: 8, 9

# Substitution Problems

Given:

$$f(a) = b$$

$$f(b) = a$$

Prove:

$$f(f(a)) = a$$

---

Given:

$$\forall x. \text{older}(\text{father}(x), x)$$

$$\text{father}(\text{bob}) = \text{art}$$

Prove:

$$\text{older}(\text{art}, x)$$

# Substitution Axioms

## Unary Relations

$$\forall x. \forall y. (p(x) \wedge x=y \Rightarrow p(y))$$

## Binary Relations

$$\forall u. \forall v. \forall x. \forall y. (q(u,v) \wedge u=x \wedge v=y \Rightarrow q(x,y))$$

## Unary Functions

$$\forall x. \forall y. \forall z. (f(x)=z \wedge x=y \Rightarrow f(y)=z)$$

## Binary Functions

$$\forall u. \forall v. \forall x. \forall y. \forall z. (g(u,v)=z \wedge u=x \wedge v=y \Rightarrow g(x,y)=z)$$



# Substitution Proof

1.  $f(a) = b$  Premise
2.  $f(b) = a$  Premise
3.  $\forall x.(x = x)$  Premise
4.  $\forall x.\forall y.(x = y \Rightarrow y = x)$  Premise
5.  $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$  Premise
6.  $\forall x.\forall y.\forall z.(f(x)=z \wedge x=y \Rightarrow f(y)=z)$  Premise

# Substitution Proof

1.  $f(a) = b$  Premise
2.  $f(b) = a$  Premise
3.  $\forall x.(x = x)$  Premise
4.  $\forall x.\forall y.(x = y \Rightarrow y = x)$  Premise
5.  $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$  Premise
6.  $\forall x.\forall y.\forall z.(f(x)=z \wedge x=y \Rightarrow f(y)=z)$  Premise
7.  $f(b)=a \wedge b=f(a) \Rightarrow f(f(a))=a$  3 x UE: 6
8.  $f(a)=b \Rightarrow b=f(a)$  2 x UE: 4
9.  $b=f(a)$  IE: 8, 1

# Substitution Proof

1.	$f(a) = b$	Premise
2.	$f(b) = a$	Premise
3.	$\forall x.(x = x)$	Premise
4.	$\forall x.\forall y.(x = y \Rightarrow y = x)$	Premise
5.	$\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$	Premise
6.	$\forall x.\forall y.\forall z.(f(x)=z \wedge x=y \Rightarrow f(y)=z)$	Premise
7.	$f(b)=a \wedge b=f(a) \Rightarrow f(f(a))=a$	3 x UE: 6
8.	$f(a)=b \Rightarrow b=f(a)$	2 x UE: 4
9.	$b=f(a)$	IE: 8, 1
10.	$f(b) = a \wedge b=f(a)$	AI: 2, 9
11.	$f(f(a))=a$	IE: 7, 10

*Too tedious. Too long.*

# Rules of Inference

Equality Introduction (QI):

$$\frac{}{\tau = \tau}$$

where  $\tau$  is a ground term

Equality Elimination (QE) - also called *paramodulation*:

$$\frac{\phi}{\sigma = \tau}}{\phi_{\sigma \leftarrow \tau}}$$

where  $\sigma$  and  $\tau$  are ground terms

NB:  $\phi_{\sigma \leftarrow \tau}$  is a copy of  $\phi$  with *0 or more* occurrences of  $\sigma$  replaced by  $\tau$ .

NB: Works using the equality in the opposite direction as well.

# Substitution Proof

1.  $f(a) = b$  Premise
2.  $f(b) = a$  Premise
3.  $f(f(a)) = a$  QE: 2, 1

# Substitution Proof

- |  |          |
|--|----------|
| 1. $\forall x. \text{older}(\text{father}(x), x)$        | Premise  |
| 2. $\text{father}(\text{bob}) = \text{art}$              | Premise  |
| 3. $\text{older}(\text{father}(\text{bob}), \text{bob})$ | UE:1     |
| 4. $\text{older}(\text{art}, \text{bob})$                | QE: 3, 2 |

# Fitch

Undo   Copy   Paste   Load   Save   Library   Help

+  - Objects:

+  - Functions:

- Select All
- 1.  $f(a)=b$  Premise
- 2.  $f(b)=a$  Premise
- 3.  $f(f(a))=a$  Equality Elimination: 1, 2

Premise	Negation Introduction	Assumption	Universal Introduction	Domain Closure
Reiteration	Negation Elimination	Implication Introduction	Universal Elimination	Induction
Truthtable	And Introduction	Implication Elimination	Existential Introduction	Equality Introduction
Shortcut	And Elimination	Biconditional Introduction	Existential Elimination	Equality Elimination
Replace	Or Introduction	Biconditional Elimination		Rational Equation
Coalesce	Or Elimination			Rational Induction
Delete				

# Fitch

Undo Copy Paste Load Save Library Help

+  - Objects:

+  - Functions:

- Select All
- 1. AX:older(father(X),X) Premise
- 2. father(bob)=art Premise
- 3. older(father(bob),bob) Universal Elimination: 1
- 4. older(art,bob) Equality Elimination: 2, 3

Premise	Negation Introduction	Assumption	Universal Introduction	Domain Closure
Reiteration	Negation Elimination	Implication Introduction	Universal Elimination	Induction
Truthtable	And Introduction	Implication Elimination	Existential Introduction	Equality Introduction
Shortcut	And Elimination	Biconditional Introduction	Existential Elimination	Equality Elimination
Replace	Or Introduction	Biconditional Elimination		Rational Equation
Coalesce	Or Elimination			Rational Induction
Delete				



# Evaluable Terms

# Signature

Object Constant: 0, ...

Unary Function Constant:  $s$

Evaluable Function Constants:  $+$ ,  $\times$


Binary Relation Constant:  $=$

*Evaluable functions do **not** expand Herbrand universe.*

*Evaluable terms **refer** to terms in Herbrand universe.*

*Evaluable functions **defined** in terms of other concepts.*

# Different Types of Terms



# Significance

For induction to work, we must show that a property holds of **all terms in the Herbrand Universe** (Herbrand terms).

When some objects or functions are defined in terms of others, it is not necessary to show that a property holds of **evaluable terms** (since all sentences involving those terms necessarily have the same truth values as sentences with equivalent **Herbrand terms**).

*Upshot: Need to do induction only on objects and functions comprising the Herbrand universe.*

# Binary Function Example

Object Constant: 0, ...

Unary Function Constant:  $s$

Binary Evaluable Function Constant:  $f$

Binary Relation Constant:  $=$

# Binary Function Problem

Axioms for +:

$$\forall y.(y + 0 = y)$$
$$\forall x.\forall y.(x + s(y) = s(x + y))$$

Problem: Prove that 0 is a left identity for +.

$$\forall y.(0 + y = y)$$

# Inductive Proof

1.  $\forall x. (x + 0 = x)$  Premise
2.  $\forall x. \forall y. (x + s(y) = s(x + y))$  Premise

# Inductive Proof

1.  $\forall x. (x + 0 = x)$  Premise
2.  $\forall x. \forall y. (x + s(y) = s(x + y))$  Premise
3.  $0 + 0 = 0$  UE: 1



# Inductive Proof

1.  $\forall x. (x + 0 = x)$  Premise
2.  $\forall x. \forall y. (x + s(y) = s(x + y))$  Premise
3.  $0 + 0 = 0$  UE: 1
4.  $| 0 + c = c$  Assumption

# Inductive Proof

- |    |   |            |
|----|---|------------|
| 1  | $\forall x. (x + 0 = x)$                          | Premise    |
| 2. | $\forall x. \forall y. (x + s(y) = s(x + y))$     | Premise    |
| 3. | $0 + 0 = 0$                                       | UE: 1      |
| 4. | $\left  0 + c = c \right.$                        | Assumption |
| 5. | $\left  \forall y. (0 + s(y) = s(0 + y)) \right.$ | UE: 2      |

# Inductive Proof

1	$\forall x. (x + 0 = x)$	Premise
2.	$\forall x. \forall y. (x + s(y) = s(x + y))$	Premise
3.	$0 + 0 = 0$	UE: 1
4.	$\left  0 + c = c \right.$	Assumption
5.	$\left  \forall y. (0 + s(y) = s(0 + y)) \right.$	UE: 2
6.	$\left  0 + s(c) = s(0 + c) \right.$	UE: 5

# Inductive Proof

1	$\forall x. (x + 0 = x)$	Premise
2.	$\forall x. \forall y. (x + s(y) = s(x + y))$	Premise
3.	$0 + 0 = 0$	UE: 1
4.	$0 + c = c$	Assumption
5.	$\forall y. (0 + s(y) = s(0 + y))$	UE: 2
6.	$0 + s(c) = s(0 + c)$	UE: 5
7.	$0 + s(c) = s(c)$	QE: 6, 4





# Inductive Proof

1	$\forall x. (x + 0 = x)$	Premise
2.	$\forall x. \forall y. (x + s(y) = s(x + y))$	Premise
3.	$0 + 0 = 0$	UE: 1
4.	$\left  0 + c = c \right.$	Assumption
5.	$\left  \forall y. (0 + s(y) = s(0 + y)) \right.$	UE: 2
6.	$\left  0 + s(c) = s(0 + c) \right.$	UE: 5
7.	$\left  0 + s(c) = s(c) \right.$	QE: 6, 4
8.	$0 + c = c \Rightarrow 0 + s(c) = s(c)$	$\Pi$ : 4, 7

# Inductive Proof

1	$\forall x. (x + 0 = x)$	Premise
2.	$\forall x. \forall y. (x + s(y) = s(x + y))$	Premise
3.	$0 + 0 = 0$	UE: 1
4.	$\left  \begin{array}{l} 0 + c = c \\ \forall y. (0 + s(y) = s(0 + y)) \\ 0 + s(c) = s(0 + c) \\ 0 + s(c) = s(c) \end{array} \right.$	Assumption
5.		UE: 2
6.		UE: 5
7.		QE: 6, 4
8.	$0 + c = c \Rightarrow 0 + s(c) = s(c)$	II: 4, 7
9.	$\forall y. (0 + y = y \Rightarrow 0 + s(y) = s(y))$	UI: 8
10.	$\forall y. (0 + y = y)$	Ind: 3, 9



# Trees



# Tree Vocabulary

Object constants:  $a, b$

Unary function constants:  $cons$





Binary relation constant:  $equal$



# Tree Vocabulary

Object constants:  $a, b$



Unary function constants:  $cons$



Unary evaluable function:  $rev$

Binary relation constant:  $equal$

# Reversing Trees



Reversing Atomic Trees:

$$\text{rev}(a) = a$$

$$\text{rev}(b) = b$$

Reversing Compound Trees:

$$\forall x. (\text{rev}(\text{cons}(x, y)) = \text{cons}(\text{rev}(y), \text{rev}(x)))$$

# Problem

Reverse is its own inverse. (In other words, the result of reversing a list twice is equal to the original list.)

$$\forall x. rev(rev(x)) = x$$

Let's prove it using induction.

Hint: *rev* is defined in terms of *cons* so we just need to do induction on *cons*.

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $\mid rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $\left| rev(rev(c)) = c \wedge rev(rev(d)) = d \right.$  Assumption
7.  $\left| rev(rev(cons(c, d))) = rev(rev(cons(c, d))) \right.$  QI

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption
7.  $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$  QI
8.  $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$  UE QE: 7, 3

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption
7.  $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$  QI
8.  $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$  UE QE: 7, 3
9.  $rev(rev(cons(c, d))) = cons(rev(rev(c)), rev(rev(d)))$  UE QE: 8, 3



# Proof By Induction

- |     |  |              |
|-----|--|--------------|
| 1.  | $rev(a) = a$   | Premise      |
| 2.  | $rev(b) = b$   | Premise      |
| 3.  | $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$ | Premise      |
| 4.  | $rev(rev(a)) = a$  | QE: 1, 1     |
| 5.  | $rev(rev(b)) = b$  | QE: 2, 2     |
| 6.  | $rev(rev(c)) = c \wedge rev(rev(d)) = d$                         | Assumption   |
| 7.  | $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$                    | QI           |
| 8.  | $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$               | UE QE: 7, 3  |
| 9.  | $rev(rev(cons(c, d))) = cons(rev(rev(c)), rev(rev(d)))$          | UE QE: 8, 3  |
| 10. | $rev(rev(cons(c, d))) = cons(c, d)$                              | 2 x QE: 9, 6 |

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption
7.  $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$  QI
8.  $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$  UE QE: 7, 3
9.  $rev(rev(cons(c, d))) = cons(rev(rev(c)), rev(rev(d)))$  UE QE: 8, 3
10.  $rev(rev(cons(c, d))) = cons(c, d)$  2 x QE: 9, 6
11.  $rev(rev(c)) = c \wedge rev(rev(d)) = d \Rightarrow rev(rev(cons(c, d))) = cons(c, d)$  II: 6, 10

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption
7.  $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$  QI
8.  $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$  UE QE: 7, 3
9.  $rev(rev(cons(c, d))) = cons(rev(rev(c)), rev(rev(d)))$  UE QE: 8, 3
10.  $rev(rev(cons(c, d))) = cons(c, d)$  2 x QE: 9, 6
11.  $rev(rev(c)) = c \wedge rev(rev(d)) = d \Rightarrow rev(rev(cons(c, d))) = cons(c, d)$  II: 6, 10
12.  $\forall x. \forall y. (rev(rev(x)) = x \wedge rev(rev(y)) = y \Rightarrow$   
 $rev(rev(cons(x, y))) = cons(x, y))$  2 x UI: 11

# Proof By Induction

1.  $rev(a) = a$  Premise
2.  $rev(b) = b$  Premise
3.  $\forall x. \forall y. (rev(cons(x, y)) = cons(rev(y), rev(x)))$  Premise
4.  $rev(rev(a)) = a$  QE: 1, 1
5.  $rev(rev(b)) = b$  QE: 2, 2
6.  $rev(rev(c)) = c \wedge rev(rev(d)) = d$  Assumption
7.  $rev(rev(cons(c, d))) = rev(rev(cons(c, d)))$  QI
8.  $rev(rev(cons(c, d))) = rev(cons(rev(d), rev(c)))$  UE QE: 7, 3
9.  $rev(rev(cons(c, d))) = cons(rev(rev(c)), rev(rev(d)))$  UE QE: 8, 3
10.  $rev(rev(cons(c, d))) = cons(c, d)$  2 x QE: 9, 6
11.  $rev(rev(c)) = c \wedge rev(rev(d)) = d \Rightarrow rev(rev(cons(c, d))) = cons(c, d)$  II: 6, 10
12.  $\forall x. \forall y. (rev(rev(x)) = x \wedge rev(rev(y)) = y \Rightarrow$   
 $rev(rev(cons(x, y))) = cons(x, y))$  2 x UI: 11
13.  $\forall x. (rev(rev(x)) = x)$  Ind: 4, 5, 12

# Polynomial Arithmetic

# Signature for Arithmetic

Object Constant: 0

Unary Function Constant:  $s$

Binary Evaluable Function Constants:

*plus* - addition

*times* - multiplication

Binary Relation Constant: =

# More Syntactic Sugar

Formal Syntax:

*equal(plus(s(0),s(0)), s(s(0)))*

*equal(times(s(s(0)),s(0)), s(s(0)))*

Syntactic Sugar:

$$1 + 1 = 2$$

$$2 \times 1 = 2$$

# Definitions of Evaluable Arithmetic Functions

Addition:

$$\forall x.(x + 0 = x)$$
$$\forall x.\forall y.(x + (y + 1) = (x + y) + 1)$$

Multiplication:

$$\forall y.(0 \times y = 0)$$
$$\forall x.\forall y.((x + 1) \times y = (x \times y) + y)$$



# Equality Proof

1.  $b = a$  Premise
2.  $b = c$  Premise
3.  $\forall x.(x = x)$  Premise
4.  $\forall x.\forall y.(x = y \Rightarrow y = x)$  Premise
5.  $\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$  Premise

# Long Messy Proofs

1.	$f(a) = b$	Premise
2.	$f(b) = a$	Premise
3.	$\forall x.(x = x)$	Premise
4.	$\forall x.\forall y.(x = y \Rightarrow y = x)$	Premise
5.	$\forall x.\forall y.\forall z.(x = y \wedge y = z \Rightarrow x = z)$	Premise
6.	$\forall x.\forall y.\forall z.(f(x)=z \wedge x=y \Rightarrow f(y)=z)$	Premise
7.	$f(b)=a \wedge b=f(a) \Rightarrow f(f(a))=a$	3 x UE: 6
8.	$f(a)=b \Rightarrow b=f(a)$	2 x UE: 4
9.	$b=f(a)$	IE: 8, 1
10.	$f(b) = a \wedge b=f(a)$	AI: 2, 9
11.	$f(f(a))=a$	IE: 7, 10

# Rational Equation

Rule of Inference (Rational Equation):

---

$$\sigma = \tau$$

where  $\sigma$  and  $\tau$  are *equivalent* polynomials

Example:

---

$$(c+1)*(c+1) = c*c+2*c+1$$

# Rational Induction

+ definable in terms of  $s$ , so we need only do induction on  $s$ .

$$\frac{\begin{array}{l} \phi[0] \\ \forall x.(\phi[x] \Rightarrow \phi[s(x)]) \end{array}}{\forall x.\phi[x]}$$

Rational Induction (since  $x + 1 = s(x)$ )

$$\frac{\begin{array}{l} \phi[0] \\ \forall x.(\phi[x] \Rightarrow \phi[x + 1]) \end{array}}{\forall x.\phi[x]}$$

# Problem

Consider the following function.

$$f(0) = 0$$
$$\forall z. (f(z+1) = f(z) + (z+1) + (z+1))$$

$f(z)$  is 2 x the sum of numbers from 0 through  $z$ .

# Problem

Function definition:

$$f(0) = 0$$
$$\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$$

**Claim:**  $f$  can be defined *non-recursively* in terms of  $+$  and  $\times$ .

$$\forall z.(f(z) = z \times (z + 1))$$

Let's prove it, using induction.

# Problem

1.  $f(0) = 0$  Premise
2.  $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$  Premise

# Problem

1.  $f(0) = 0$  Premise
2.  $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$  Premise
3.  $0 \times (0 + 1) = 0$  Equation



# Problem

- |    |   |          |
|----|---|----------|
| 1. | $f(0) = 0$                                  | Premise  |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$ | Premise  |
| 3. | $0 = 0 \times (0 + 1)$                      | Equation |
| 4. | $f(0) = 0 \times (0 + 1)$                   | QE: 3, 1 |

Base Case for :

$$\forall z.(f(z) = z \times (z + 1))$$

# Problem

- |    |   |            |
|----|---|------------|
| 1  | $f(0) = 0$                                  | Premise    |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$ | Premise    |
| 3. | $0 \times (0 + 1) = 0$                      | Equation   |
| 4. | $f(0) = 0 \times (0 + 1)$                   | QE: 3, 1   |
| 5. | $\mid f(c) = c \times (c+1)$                | Assumption |

We want to prove:

$$\forall z.(f(z) = z \times (z + 1) \Rightarrow f(z+1) = (z + 1) \times ((z + 1) + 1))$$

To do that, we need to prove:

$$f(c) = c \times (c + 1) \Rightarrow f(c+1) = (c + 1) \times ((c + 1) + 1)$$

To do that, we assume  $f(c) = c \times (c + 1)$

and prove  $f(c+1) = (c + 1) \times ((c + 1) + 1)$

# Problem

- |    |  |            |
|----|--|------------|
| 1  | $f(0) = 0$   | Premise    |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$  | Premise    |
| 3. | $0 \times (0 + 1) = 0$   | Equation   |
| 4. | $f(0) = 0 \times (0 + 1)$  | QE: 3, 1   |
| 5. | $\left  \begin{array}{l} f(c) = c \times (c+1) \\ f(c+1) = f(c) + (c+1) + (c+1) \end{array} \right.$ | Assumption |
| 6. |  | UE: 2      |

# Problem

- |    |   |            |
|----|---|------------|
| 1. | $f(0) = 0$                                  | Premise    |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$ | Premise    |
| 3. | $0 \times (0 + 1) = 0$                      | Equation   |
| 4. | $f(0) = 0 \times (0 + 1)$                   | QE: 3, 1   |
| 5. | $f(c) = c \times (c+1)$                     | Assumption |
| 6. | $f(c+1) = f(c) + (c+1) + (c+1)$             | UE: 2      |
| 7. | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$   | QE: 5, 6   |

# Problem

- |    |   |            |
|----|---|------------|
| 1. | $f(0) = 0$  | Premise    |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$                 | Premise    |
| 3. | $0 \times (0 + 1) = 0$                                      | Equation   |
| 4. | $f(0) = 0 \times (0 + 1)$                                   | QE: 3, 1   |
| 5. | $f(c) = c \times (c+1)$                                     | Assumption |
| 6. | $f(c+1) = f(c) + (c+1) + (c+1)$                             | UE: 2      |
| 7. | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$                   | QE: 5, 6   |
| 8. | $c \times (c+1) + (c+1) + (c+1) = (c+1) \times ((c+1) + 1)$ | Equation   |

# Problem

- |    |   |            |
|----|---|------------|
| 1. | $f(0) = 0$  | Premise    |
| 2. | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$                 | Premise    |
| 3. | $0 \times (0 + 1) = 0$                                      | Equation   |
| 4. | $f(0) = 0 \times (0 + 1)$                                   | QE: 3, 1   |
| 5. | $f(c) = c \times (c+1)$                                     | Assumption |
| 6. | $f(c+1) = f(c) + (c+1) + (c+1)$                             | UE: 2      |
| 7. | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$                   | QE: 5, 6   |
| 8. | $c \times (c+1) + (c+1) + (c+1) = (c+1) \times ((c+1) + 1)$ | Equation   |
| 9. | $f(c+1) = (c+1) \times ((c+1) + 1)$                         | QE: 8, 7   |

# Problem

- |     |   |            |
|-----|---|------------|
| 1.  | $f(0) = 0$  | Premise    |
| 2.  | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$                           | Premise    |
| 3.  | $0 \times (0 + 1) = 0$  | Equation   |
| 4.  | $f(0) = 0 \times (0 + 1)$   | QE: 3, 1   |
| 5.  | $f(c) = c \times (c+1)$   | Assumption |
| 6.  | $f(c+1) = f(c) + (c+1) + (c+1)$                                       | UE: 2      |
| 7.  | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$                             | QE: 5, 6   |
| 8.  | $c \times (c+1) + (c+1) + (c+1) = (c+1) \times ((c+1) + 1)$           | Equation   |
| 9.  | $f(c+1) = (c+1) \times ((c+1) + 1)$                                   | QE: 8, 7   |
| 10. | $f(c) = c \times (c+1) \Rightarrow f(c+1) = (c+1) \times ((c+1) + 1)$ | II: 5, 9   |

# Problem

- |     |   |            |
|-----|---|------------|
| 1.  | $f(0) = 0$  | Premise    |
| 2.  | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$                                       | Premise    |
| 3.  | $0 \times (0 + 1) = 0$  | Equation   |
| 4.  | $f(0) = 0 \times (0 + 1)$   | QE: 3, 1   |
| 5.  | $f(c) = c \times (c+1)$   | Assumption |
| 6.  | $f(c+1) = f(c) + (c+1) + (c+1)$   | UE: 2      |
| 7.  | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$   | QE: 5, 6   |
| 8.  | $c \times (c+1) + (c+1) + (c+1) = (c+1) \times ((c+1) + 1)$                       | Equation   |
| 9.  | $f(c+1) = (c+1) \times ((c+1) + 1)$   | QE: 8, 7   |
| 10. | $f(c) = c \times (c+1) \Rightarrow f(c+1) = (c+1) \times ((c+1) + 1)$             | II: 5, 9   |
| 11. | $\forall z.(f(z) = z \times (z+1) \Rightarrow f(z+1) = (z+1) \times ((z+1) + 1))$ | UI: 10     |



# Problem

- |     |   |               |
|-----|---|---------------|
| 1.  | $f(0) = 0$  | Premise       |
| 2.  | $\forall z.(f(z+1) = f(z) + (z+1) + (z+1))$                                       | Premise       |
| 3.  | $0 \times (0 + 1) = 0$  | Equation      |
| 4.  | $f(0) = 0 \times (0 + 1)$   | QE: 3, 1      |
| 5.  | $f(c) = c \times (c+1)$   | Assumption    |
| 6.  | $f(c+1) = f(c) + (c+1) + (c+1)$   | UE: 2         |
| 7.  | $f(c+1) = c \times (c+1) + (c+1) + (c+1)$   | QE: 5, 6      |
| 8.  | $c \times (c+1) + (c+1) + (c+1) = (c+1) \times ((c+1) + 1)$                       | Equation      |
| 9.  | $f(c+1) = (c+1) \times ((c+1) + 1)$   | QE: 8, 7      |
| 10. | $f(c) = c \times (c+1) \Rightarrow f(c+1) = (c+1) \times ((c+1) + 1)$             | II: 5, 9      |
| 11. | $\forall z.(f(z) = z \times (z+1) \Rightarrow f(z+1) = (z+1) \times ((z+1) + 1))$ | UI: 10        |
| 12. | $\forall z.(f(z) = z \times (z+1))$   | Ratind: 4, 11 |

# Fitch

Undo   Copy   Paste   Load   Save   Help

+  - Objects:

+  - Functions:

<input type="checkbox"/>	Select All	
<input type="checkbox"/> 1.	$f(0)=0$	Premise
<input type="checkbox"/> 2.	$AZ:f(Z+1)=f(Z)+((Z+1)+(Z+1))$	Premise
<input type="checkbox"/> 3.	$0*(0+1)=0$	Equation
<input type="checkbox"/> 4.	$f(0)=0*(0+1)$	Equality Elimination: 3, 1
<input type="checkbox"/> 5.	$f(c)=c*(c+1)$	Assumption
<input type="checkbox"/> 6.	$f(c+1)=f(c)+((c+1)+(c+1))$	Universal Elimination: 2
<input type="checkbox"/> 7.	$f(c+1)=c*(c+1)+((c+1)+(c+1))$	Equality Elimination: 5, 6
<input type="checkbox"/> 8.	$c*(c+1)+((c+1)+(c+1))=(c+1)*((c+1)+1)$	Equation
<input type="checkbox"/> 9.	$f(c+1)=(c+1)*((c+1)+1)$	Equality Elimination: 8, 7
<input type="checkbox"/> 10.	$f(c)=c*(c+1) \Rightarrow f(c+1)=(c+1)*((c+1)+1)$	Implication Introduction: 5, 9
<input type="checkbox"/> 11.	$AZ:(f(Z)=Z*(Z+1) \Rightarrow f(Z+1)=(Z+1)*((Z+1)+1))$	Universal Introduction: 10
<input type="checkbox"/> 12.	$AZ:f(Z)=Z*(Z+1)$	Rational Induction
Goal	$AZ:f(Z)=Z*(Z+1)$	Complete

# Yet Another Problem

Recursive Function:

$$g(0) = 0$$

$$g(z + 1) = g(z) + (2 \times z + 1)$$

Non-recursive definition in terms of  $\times$ :

$$g(z) = z \times z$$


$z$	$g(z)$
0	0
1	1
2	4
3	9

# And Another

Number of cans per layer (starting at 1)

$$h(1) = 1$$

$$h(z + 1) = h(z) + 6 \times z$$



Non-recursive definition in terms of + and  $\times$ :

$$h(z+1) = 3 \times z \times (z + 1) + 1$$

